

ELLIOT M. HIRSCH, MD PLASTIC & RECONSTRUCTIVE SURGERY

4955 Van Nuys Blvd, Suite 715, Sherman Oaks, California, 91403 Phone: 818-825-8131 • Fax: 818-616-1044 • www.hirschplasticsurgery.com

Nipple Reconstruction Post-Operative Instructions

<u>Medications</u>: Your greatest discomfort will be in the first 24-48 hours after your procedure. If you have pain, you should take your prescribed pain medication or Tylenol. After 48-72 hours, you can take Ibuprofen if desired. Take your pain medication with food to avoid nausea or stomach discomfort. Pain medication can be constipating, so be sure to eat fruits and vegetables. Drink plenty of fluids as well.

<u>Dressings and Incision Care</u>: Leave your dressing in place for 48 hours, then it is ok to remove it and shower. Wear a soft, loose fitting bra after you remove your dressing, but do not wear any compressive wraps or garments.

<u>Activity</u>: Make sure that you are up and walking and moving around the first night after surgery, this is very important! You will find that the more you move around, the better you feel. Please avoid strenuous activity or any lifting greater than 20 pounds until you are seen by Dr. Hirsch.

<u>Diet</u> : You may resume your regular diet
Special Instructions:
When to Call: A small to moderate amount of clear to light pink drainage from the incision is normal. However, if this drainage is dark, bloody, or foul smelling, please call Dr. Hirsch.

- fever of 101 or higher
- severe pain not responsive to pain medications

Please call Dr. Hirsch's office if you are experiencing any of the following:

- increasing swelling
- increasing redness
- pus or foul-smelling drainage from the incision.

If you have any other questions or concerns, feel free to call our office at any time: 818-825	-8131.
Follow up:	